



Spring Linseed Bulletin

06/15

Linseed Harvesting

Linseed crops are turning quickly, meaning desiccation is near. For most growers the Linseed will be harvested straight after their Wheat. We are expecting the early crops to be harvested in the third week of August.

When to Harvest

Typically, crops should be ready for harvest 5 – 10 days after the application of Diquat

- The top half of the stem will be dead, although the lower half is still likely to be yellowy/green.
- Capsules will be 'parchment' colour.
- Seeds will be dark brown and will easily rub out from the capsules.
- Moisture content of the seed will be 8.5 – 12%.

When the Linseed stem is at the stage shown in the picture, the fibres (which are arranged in bundles on the outside of the stem) are tightly 'glued' to the inner woody core, making the straw easy to cut.

As the stem ages, this 'glue' gradually breaks down – a natural process known as retting. This allows the fibres to separate from the stem easily. This means that:

- a) the stem becomes more difficult to cut
- b) it is more likely to wrap in the combine
- c) the straw is less likely to chop



Linseed stem with fibre peeled back, exposing the inner woody core

Cut at the first opportunity and do not leave Linseed too long after you have finished your Wheat harvest. Whilst Linseed does not shed easily, leaving Linseed too long will result in the stems "retting" and becoming more difficult to cut and more likely to wrap.

Combines will cut green Linseed straw provided the knife and fingers are in good condition.

- Use seed maturity/moisture content as your combining guide.
- Linseed cuts more easily nearer the ground (see Combine Operator Instructions).
- Do not miss any opportunity to combine Linseed.
- Your combine is the best judge of whether to harvest Linseed!
- You will often find with Linseed that one day harvesting will not 'go', but the next day for no apparent reason you will fly through the crop.

Combine Settings

Outline instructions can be downloaded from:

www.premiumcrops.com/usefuldocuments.

Market Standards

Normal trading standards for Linseed are:

- Moisture basis 9%, maximum 10%
- Admixture 2%, maximum 4%
- Mite free
- Clover weevil free



Linseed Straw Disposal

There are several options for Linseed straw disposal:

Chopping - In some years Linseed straw will chop, but the combine chopper blades need to be in tip-top condition.

Problems can arise when the straw is weathered, which results in a fluffy material which can hinder subsequent min-till operations. Our advice is to disengage the chopper if you are not obtaining a 'clean chop'.

Baling - Some growers use Linseed straw for sugarbeet clamps or for bedding.

Burning - Burning is the simplest and most widely used method of straw disposal. Linseed straw is exempt from the burning ban and burns quickly without smoke or smuts. Either push the straw into a line across the centre of the field or round bale without strings.

Burning must be carried out in accordance with the appropriate legislation (The Crop Residues (Burning) Regulations 1993).

Drying and Storage

- Most Linseed will be harvested in the 8.0 – 12.0% moisture range – it needs to be dried to 8.0% moisture as it is going to be stored for a number of months prior to processing.
- Damp Linseed will deteriorate rapidly above 11.0% moisture.

If drying is required:

- Maximum temperature should not exceed 55°C.
 - Seed harvested at >11% may need 2 passes through the dryer
 - Maximum 1 metre depth for on-floor drying, because Linseed has a very high resistance to air-flow.
 - Take care when storing/drying damp Linseed in a bin system. The airflow resistance can cause the drying front to stall, causing wet Linseed to 'set'. If possible rotate from one bin to another.
 - Take down to 8% moisture, to allow for variations in meter readings.
-
- If harvesting in hot conditions, ensure that stored Linseed is cooled to <20°C (15°C or below ideally) when cooler weather conditions allow. This will help prevent mite infestations.
 - In some years, clover weevils are harvested with the Linseed. The weevil will dissipate over a few weeks.
 - Monitor stored linseed regularly for moisture and temperature increases, as well as mites. The easiest way to check for mites is to put a sample of your Linseed in a plastic bag, leave the bag on the window-sill and if any mites are present they will migrate to the top corners of the bag. The mites are small (<1 mm) and pinkish coloured. The only proven way to remove mites is to re-dry and clean the crop. End users are becoming more aware of mite contamination and may reject loads if found at the delivery point.
 - When Linseed is dry and cool (<8.5% moisture and <15°C), it will store quite happily for several months.

Samples

Once your crop has been dried please send us a sample (approximately 0.5 – 0.75 kg):

- Put the sample in the polythene sample bag
- Complete your trading name and estimated tonnage on the bag
- Put into the pre-addressed survivor envelope provided and post it (please make sure you get the parcel weighed at the Post Office to ensure the correct postage is paid)

We will analyse the sample and send you an analysis report in due course.

7th August 2015

Always read the label before using pesticides. Use pesticides safely.